


INSTRUCTOR'S
SOLUTIONS MANUAL

AN INTRODUCTION TO
MATHEMATICAL STATISTICS
AND ITS APPLICATIONS
SIXTH EDITION

Richard Larsen
Vanderbilt University

Morris Marx
University of West Florida





This work is protected by United States copyright laws and is provided solely for the use of instructors in teaching their courses and assessing student learning. Dissemination or sale of any part of this work (including on the World Wide Web) will destroy the integrity of the work and is not permitted. The work and materials from it should never be made available to students except by instructors using the accompanying text in their classes. All recipients of this work are expected to abide by these restrictions and to honor the intended pedagogical purposes and the needs of other instructors who rely on these materials.

The author and publisher of this book have used their best efforts in preparing this book. These efforts include the development, research, and testing of the theories and programs to determine their effectiveness. The author and publisher make no warranty of any kind, expressed or implied, with regard to these programs or the documentation contained in this book. The author and publisher shall not be liable in any event for incidental or consequential damages in connection with, or arising out of, the furnishing, performance, or use of these programs.

Reproduced by Pearson from electronic files supplied by the author.

Copyright © 2018, 2012, 2006 Pearson Education, Inc.
Publishing as Pearson, 330 Hudson Street, NY NY 10013

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior written permission of the publisher. Printed in the United States of America.



ISBN-13: 978-0-13-411427-9

ISBN-10: 0-13-411427-2

Contents

Chapter 2: Probability	1
2.2 Samples Spaces and the Algebra of Sets	1
2.3 The Probability Function	6
2.4 Conditional Probability	7
2.5 Independence	13
2.6 Combinatorics	17
2.7 Combinatorial Probability	24
Chapter 3: Random Variables	27
3.2 Binomial and Hypergeometric Probabilities	27
3.3 Discrete Random Variables	36
3.4 Continuous Random Variables	41
3.5 Expected Values	44
3.6 The Variance	52
3.7 Joint Densities	57
3.8 Transforming and Combining Random Variables	69
3.9 Further Properties of the Mean and Variance	73
3.10 Order Statistics	79
3.11 Conditional Densities	82
3.12 Moment-Generating Functions	88
Chapter 4: Special Distributions	93
4.2 The Poisson Distribution	93
4.3 The Normal Distribution	98
4.4 The Geometric Distribution	105
4.5 The Negative Binomial Distribution	107
4.6 The Gamma Distribution	109
Chapter 5: Estimation	111
5.2 Estimating Parameters: The Method of Maximum Likelihood and Method of Moments	111
5.3 Interval Estimation	118
5.4 Properties of Estimators	123
5.5 Minimum-Variance Estimators: The Cramér-Rao Lower Bound	128
5.6 Sufficient Estimators	130
5.7 Consistency	133
5.8 Bayesian Estimation	135

Chapter 6: Hypothesis Testing	137
6.2 The Decision Rule.....	137
6.3 Testing Binomial Data.....	138
6.4 Type I and Type II Errors.....	139
6.5 A Notion of Optimality: The Generalized Likelihood Ratio.....	144
Chapter 7: Inferences Based on the Normal Distribution	147
7.3 Deriving the Distribution of $\frac{\bar{Y}-\mu}{S/\sqrt{n}}$	147
7.4 Drawing Inferences About μ	150
7.5 Drawing Inferences About σ^2	156
Chapter 8: Types of Data: A Brief Overview	161
8.2 Classifying Data.....	161
Chapter 9: Two-Sample Inference	163
9.2 Testing $H_0 : \mu_X = \mu_Y$	163
9.3 Testing $H_0 : \sigma_X^2 = \sigma_Y^2$ —The F Test.....	166
9.4 Binomial Data: Testing $H_0 : p_X = p_Y$	168
9.5 Confidence Intervals for the Two-Sample Problem.....	170
Chapter 10: Goodness-of-Fit Tests	173
10.2 The Multinomial Distribution.....	173
10.3 Goodness-of-Fit Tests: All Parameters Known.....	175
10.4 Goodness-of-Fit Tests: Parameters Unknown.....	178
10.5 Contingency Tables.....	185
Chapter 11: Regression	189
11.2 The Method of Least Squares.....	189
11.3 The Linear Model.....	199
11.4 Covariance and Correlation.....	204
11.5 The Bivariate Normal Distribution.....	208
Chapter 12: The Analysis of Variance	211
12.2 The F test.....	211
12.3 Multiple Comparisons: Tukey’s Method.....	214

12.4	Testing Subhypotheses with Constrasts	216
12.5	Data Transformations	218
Appendix 12.A.2	The Distribution of $\frac{SSTR/(k-1)}{SSE/(n-k)}$ When H_1 Is True	218
Chapter 13: Randomized Block Designs		221
13.2	The F Test for a Randomized Block Design	221
13.3	The Paired t Test.....	224
Chapter 14: Nonparametric Statistics.....		229
14.2	The Sign Test	229
14.3	Wilcoxon Tests.....	232
14.4	The Kruskal-Wallis Test	236
14.5	The Friedman Test.....	239
14.6	Testing for Randomness.....	241

Chapter 2: Probability


Section 2.2: Sample Spaces and the Algebra of Sets

- 2.2.1 $S = \{(s,s,s), (s,s,f), (s,f,s), (f,s,s), (s,f,f), (f,s,f), (f,f,s), (f,f,f)\}$
 $A = \{(s,f,s), (f,s,s)\}; B = \{(f,f,f)\}$
- 2.2.2 Let (x, y, z) denote a red x , a blue y , and a green z .
Then $A = \{(2,2,1), (2,1,2), (1,2,2), (1,1,3), (1,3,1), (3,1,1)\}$
- 2.2.3 $(1,3,4), (1,3,5), (1,3,6), (2,3,4), (2,3,5), (2,3,6)$
- 2.2.4 There are 16 ways to get an ace and a 7, 16 ways to get a 2 and a 6, 16 ways to get a 3 and a 5, and 6 ways to get two 4's, giving 54 total.
- 2.2.5 The outcome sought is $(4, 4)$. It is "harder" to obtain than the set $\{(5, 3), (3, 5), (6, 2), (2, 6)\}$ of other outcomes making a total of 8.
- 2.2.6 The set N of five card hands in hearts that are not flushes are called *straight flushes*. These are five cards whose denominations are consecutive. Each one is characterized by the lowest value in the hand. The choices for the lowest value are A, 2, 3, ..., 10. (Notice that an ace can be high or low). Thus, N has 10 elements.
- 2.2.7 $P = \{\text{right triangles with sides } (5, a, b): a^2 + b^2 = 25\}$
- 2.2.8 $A = \{SSBBBB, SBSBBB, SBBSBB, SBBBSB, BSSBBB, BSBSBB, BSBBBS, BBSSBB, BBSBSB, BBBSSB\}$
- 2.2.9 (a) $S = \{(0, 0, 0, 0), (0, 0, 0, 1), (0, 0, 1, 0), (0, 0, 1, 1), (0, 1, 0, 0), (0, 1, 0, 1), (0, 1, 1, 0), (0, 1, 1, 1), (1, 0, 0, 0), (1, 0, 0, 1), (1, 0, 1, 0), (1, 0, 1, 1), (1, 1, 0, 0), (1, 1, 0, 1), (1, 1, 1, 0), (1, 1, 1, 1)\}$
(b) $A = \{(0, 0, 1, 1), (0, 1, 0, 1), (0, 1, 1, 0), (1, 0, 0, 1), (1, 0, 1, 0), (1, 1, 0, 0)\}$
(c) $1 + k$
- 2.2.10 (a) $S = \{(1, 1), (1, 2), (1, 4), (2, 1), (2, 2), (2, 4), (4, 1), (4, 2), (4, 4)\}$
(b) $\{2, 3, 4, 5, 6, 8\}$
- 2.2.11 Let p_1 and p_2 denote the two perpetrators and $i_1, i_2,$ and i_3 , the three in the lineup who are innocent.
Then $S = \{(p_1, i_1), (p_1, i_2), (p_1, i_3), (p_2, i_1), (p_2, i_2), (p_2, i_3), (p_1, p_2), (i_1, i_2), (i_1, i_3), (i_2, i_3)\}$.
The event A contains every outcome in S except (p_1, p_2) .
- 2.2.12 The quadratic equation will have complex roots—that is, the event A will occur—if $b^2 - 4ac < 0$.
- 2.2.13 In order for the shooter to win with a point of 9, one of the following (countably infinite) sequences of sums must be rolled: $(9,9), (9, \text{no } 7 \text{ or no } 9,9), (9, \text{no } 7 \text{ or no } 9, \text{no } 7 \text{ or no } 9,9), \dots$

2.2.14 Let (x, y) denote the strategy of putting x white chips and y red chips in the first urn (which results in $10 - x$ white chips and $10 - y$ red chips being in the second urn). Then $S = \{(x, y) : x = 0, 1, \dots, 10, y = 0, 1, \dots, 10, \text{ and } 1 \leq x + y \leq 19\}$. Intuitively, the optimal strategies are $(1, 0)$ and $(9, 10)$.

2.2.15 Let A_k be the set of chips put in the urn at $1/2^k$ minute until midnight. For example, $A_1 = \{11, 12, 13, 14, 15, 16, 17, 18, 19, 20\}$. Then the set of chips in the urn at midnight is $\bigcup_{k=1}^{\infty} (A_k - \{k+1\}) = \emptyset$.


2.2.16 move arrow on first figure raise B by 1



2.2.17 If $x^2 + 2x \leq 8$, then $(x + 4)(x - 2) \leq 0$ and $A = \{x : -4 \leq x \leq 2\}$. Similarly, if $x^2 + x \leq 6$, then $(x + 3)(x - 2) \leq 0$ and $B = \{x : -3 \leq x \leq 2\}$. Therefore, $A \cap B = \{x : -3 \leq x \leq 2\}$ and $A \cup B = \{x : -4 \leq x \leq 2\}$.

2.2.18 $A \cap B \cap C = \{x : x = 2, 3, 4\}$

2.2.19 The system fails if either the first pair fails or the second pair fails (or both pairs fail). For either pair to fail, though, both of its components must fail. Therefore, $A = (A_{11} \cap A_{21}) \cup (A_{12} \cap A_{22})$.



2.2.21 40

2.2.22 (a) $\{E1, E2\}$ (b) $\{S1, S2, T1, T2\}$ (c) $\{A, I\}$

2.2.23 (a) If s is a member of $A \cup (B \cap C)$ then s belongs to A or to $B \cap C$. If it is a member of A or of $B \cap C$, then it belongs to $A \cup B$ and to $A \cup C$. Thus, it is a member of $(A \cup B) \cap (A \cup C)$.
Conversely, choose s in $(A \cup B) \cap (A \cup C)$. If it belongs to A , then it belongs to $A \cup (B \cap C)$. If it does not belong to A , then it must be a member of $B \cap C$. In that case it also is a member of $A \cup (B \cap C)$.

(b) If s is a member of $A \cap (B \cup C)$ then s belongs to A and to $B \cup C$. If it is a member of B , then it belongs to $A \cap B$ and, hence, $(A \cap B) \cup (A \cap C)$. Similarly, if it belongs to C , it is a member of $(A \cap B) \cup (A \cap C)$. Conversely, choose s in $(A \cap B) \cup (A \cap C)$. Then it belongs to A . If it is a member of $A \cap B$ then it belongs to $A \cap (B \cup C)$. Similarly, if it belongs to $A \cap C$, then it must be a member of $A \cap (B \cup C)$.

2.2.24 Let $B = A_1 \cup A_2 \cup \dots \cup A_k$. Then $A_1^c \cap A_2^c \cap \dots \cap A_k^c = (A_1 \cup A_2 \cup \dots \cup A_k)^c = B^c$. Then the expression is simply $B \cup B^c = S$.

2.2.25 (a) Let s be a member of $A \cup (B \cap C)$. Then s belongs to either A or $B \cap C$ (or both). If s belongs to A , it necessarily belongs to $(A \cup B) \cap C$. If s belongs to $B \cap C$, it belongs to B or C or both, so it must belong to $(A \cup B) \cap C$. Now, suppose s belongs to $(A \cup B) \cap C$. Then it belongs to either $A \cup B$ or C or both. If it belongs to C , it must belong to $A \cup (B \cap C)$. If it belongs to $A \cup B$, it must belong to either A or B or both, so it must belong to $A \cup (B \cap C)$.

(b) Suppose s belongs to $A \cap (B \cap C)$, so it is a member of A and also $B \cap C$. Then it is a member of A and of B and C . That makes it a member of $(A \cap B) \cap C$. Conversely, if s is a member of $(A \cap B) \cap C$, a similar argument shows it belongs to $A \cap (B \cap C)$.

2.2.26 (a) $A^c \cap B^c \cap C^c$
 (b) $A \cap B \cap C$
 (c) $A \cap B^c \cap C^c$
 (d) $(A \cap B^c \cap C^c) \cup (A^c \cap B \cap C^c) \cup (A^c \cap B^c \cap C)$
 (e) $(A \cap B \cap C^c) \cup (A \cap B^c \cap C) \cup (A^c \cap B \cap C)$

2.2.27 A is a subset of B .

2.2.28 (a) $\{0\} \cup \{x: 5 \leq x \leq 10\}$
 (b) $\{x: 3 \leq x < 5\}$
 (c) $\{x: 0 < x \leq 7\}$
 (d) $\{x: 0 < x < 3\}$
 (e) $\{x: 3 \leq x \leq 10\}$
 (f) $\{x: 7 < x \leq 10\}$

2.2.29 (a) B and C
 (b) B is a subset of A .


2.2.30 (a) $A_1 \cap A_2 \cap A_3$
 (b) $A_1 \cup A_2 \cup A_3$

The second protocol would be better if speed of approval matters. For very important issues, the first protocol is superior.


2.2.31 Let A and B denote the students who saw the movie the first time and the second time, respectively. Then $N(A) = 850$, $N(B) = 690$, and $N[(A \cup B)^c] = 4700$ (implying that $N(A \cup B) = 1300$). Therefore, $N(A \cap B) =$ number who saw movie twice $= 850 + 690 - 1300 = 240$.

2.2.32


(a)




$$(A \cap B)^c = A^c \cup B^c$$



(b)




$$(A \cup B)^c = A^c \cap B^c$$




2.2.33


(a)




$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$



(b)




$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$




2.2.34

(a)




$$A \cup (B \cup C)$$




$$(A \cap B) \cup C$$

(b)



$$A \cap (B \cap C)$$




$$(A \cap B) \cap C$$


2.2.35

A and B are subsets of $A \cup B$.


2.2.36



$$(A \cap B^c)^c = A^c \cup B$$



$$B \cup (A \cup B)^c = A^c \cup B$$




$$A \cap (A \cap B)^c = A \cap B^c$$

2.2.37

Let A be the set of those with MCAT scores ≥ 27 and B be the set of those with GPAs ≥ 3.5 . We are given that $N(A) = 1000$, $N(B) = 400$, and $N(A \cap B) = 300$. Then $N(A^c \cap B^c) = N[(A \cup B)^c] = 1200 - N(A \cup B) = 1200 - [(N(A) + N(B) - N(A \cap B))] = 1200 - [(1000 + 400 - 300)] = 100$. The requested proportion is $100/1200$.

2.2.38



$$N(A \cup B \cup C) = N(A) + N(B) + N(C) - N(A \cap B) - N(A \cap C) - N(B \cap C) + N(A \cap B \cap C)$$

2.2.39

Let A be the set of those saying “yes” to the first question and B be the set of those saying “yes” to the second question. We are given that $N(A) = 600$, $N(B) = 400$, and $N(A^c \cap B) = 300$. Then $N(A \cap B) = N(B) - N(A^c \cap B) = 400 - 300 = 100$. $N(A \cap B^c) = N(A) - N(A \cap B) = 600 - 100 = 500$.

2.2.40

$$\begin{aligned} N[(A \cap B)^c] &= 120 - N(A \cup B) = 120 - [N(A^c \cap B) + N(A \cap B^c) + N(A \cap B)] \\ &= 120 - [50 + 15 + 2] = 53 \end{aligned}$$

Section 2.3: The Probability Function


2.3.1 Let L and V denote the sets of programs with offensive language and too much violence, respectively. Then $P(L) = 0.42$, $P(V) = 0.27$, and $P(L \cap V) = 0.10$.
Therefore, $P(\text{program complies}) = P((L \cup V)^c) = 1 - [P(L) + P(V) - P(L \cap V)] = 0.41$.

2.3.2 $P(A \text{ or } B \text{ but not both}) = P(A \cup B) - P(A \cap B) = P(A) + P(B) - P(A \cap B) - P(A \cap B) = 0.4 + 0.5 - 0.1 - 0.1 = 0.7$


2.3.3 (a) $1 - P(A \cap B)$
(b) $P(B) - P(A \cap B)$

2.3.4 $P(A \cup B) = P(A) + P(B) - P(A \cap B) = 0.3$; $P(A) - P(A \cap B) = 0.1$. Therefore, $P(B) = 0.2$.



2.3.5 No. $P(A_1 \cup A_2 \cup A_3) = P(\text{at least one "6" appears}) = 1 - P(\text{no 6's appear}) = 1 - \left(\frac{5}{6}\right)^3 \neq \frac{1}{2}$.
The A_i 's are not mutually exclusive, so $P(A_1 \cup A_2 \cup A_3) \neq P(A_1) + P(A_2) + P(A_3)$.



$$P(A \text{ or } B \text{ but not both}) = 0.5 - 0.2 = 0.3$$



By inspection, $B = (B \cap A_1) \cup (B \cap A_2) \cup \dots \cup (B \cap A_n)$.



- 2.3.9** $P(\text{odd man out}) = 1 - P(\text{no odd man out}) = 1 - P(HHH \text{ or } TTT) = 1 - \frac{2}{8} = \frac{3}{4}$
- 2.3.10** $A = \{2, 4, 6, \dots, 24\}$; $B = \{3, 6, 9, \dots, 24\}$; $A \cap B = \{6, 12, 18, 24\}$.
Therefore, $P(A \cup B) = P(A) + P(B) - P(A \cap B) = \frac{12}{24} + \frac{8}{24} - \frac{4}{24} = \frac{16}{24}$.
- 2.3.11** Let A : State wins Saturday and B : State wins next Saturday. Then $P(A) = 0.10$, $P(B) = 0.30$, and $P(\text{lose both}) = 0.65 = 1 - P(A \cup B)$, which implies that $P(A \cup B) = 0.35$. Therefore, $P(A \cap B) = 0.10 + 0.30 - 0.35 = 0.05$, so $P(\text{State wins exactly once}) = P(A \cup B) - P(A \cap B) = 0.35 - 0.05 = 0.30$.
- 2.3.12** Since A_1 and A_2 are mutually exclusive and cover the entire sample space, $p_1 + p_2 = 1$.
But $3p_1 - p_2 = \frac{1}{2}$, so $p_2 = \frac{5}{8}$.
- 2.3.13** Let F : female is hired and T : minority is hired. Then $P(F) = 0.60$, $P(T) = 0.30$, and $P(F^c \cap T^c) = 0.25 = 1 - P(F \cup T)$. Since $P(F \cup T) = 0.75$, $P(F \cap T) = 0.60 + 0.30 - 0.75 = 0.15$.
- 2.3.14** The smallest value of $P[(A \cup B \cup C)^c]$ occurs when $P(A \cup B \cup C)$ is as large as possible. This, in turn, occurs when A , B , and C are mutually disjoint. The largest value for $P(A \cup B \cup C)$ is $P(A) + P(B) + P(C) = 0.2 + 0.1 + 0.3 = 0.6$. Thus, the smallest value for $P[(A \cup B \cup C)^c]$ is 0.4.
- 2.3.15** (a) $X^c \cap Y = \{(H, T, T, H), (T, H, H, T)\}$, so $P(X^c \cap Y) = 2/16$
(b) $X \cap Y^c = \{(H, T, T, T), (T, T, T, H), (T, H, H, H), (H, H, H, T)\}$ so $P(X \cap Y^c) = 4/16$
- 2.3.16** $A = \{(1, 5), (2, 4), (3, 3), (4, 2), (5, 1)\}$
 $A \cap B^c = \{(1, 5), (3, 3), (5, 1)\}$, so $P(A \cap B^c) = 3/36 = 1/12$.
- 2.3.17** $A \cap B, (A \cap B) \cup (A \cap C), A, A \cup B, S$
- 2.3.18** Let A be the event of getting arrested for the first scam; B , for the second. We are given $P(A) = 1/10$, $P(B) = 1/30$, and $P(A \cap B) = 0.0025$. Her chances of not getting arrested are $P[(A \cup B)^c] = 1 - P(A \cup B) = 1 - [P(A) + P(B) - P(A \cap B)] = 1 - [1/10 + 1/30 - 0.0025] = 0.869$

Section 2.4: Conditional Probability

2.4.1
$$P(\text{sum} = 10 | \text{sum exceeds } 8) = \frac{P(\text{sum} = 10 \text{ and sum exceeds } 8)}{P(\text{sum exceeds } 8)}$$

$$= \frac{P(\text{sum} = 10)}{P(\text{sum} = 9, 10, 11, \text{ or } 12)} = \frac{3/36}{4/36 + 3/36 + 2/36 + 1/36} = \frac{3}{10}$$

2.4.2 $P(A|B) + P(B|A) = 0.75 = \frac{P(A \cap B)}{P(B)} + \frac{P(A \cap B)}{P(A)} = \frac{10P(A \cap B)}{4} + 5P(A \cap B)$, which implies that $P(A \cap B) = 0.1$.

2.4.3 If $P(A|B) = \frac{P(A \cap B)}{P(B)} < P(A)$, then $P(A \cap B) < P(A) \cdot P(B)$.

It follows that $P(B|A) = \frac{P(A \cap B)}{P(A)} < \frac{P(A) \cdot P(B)}{P(A)} = P(B)$.

2.4.4 $P(E|A \cup B) = \frac{P(E \cap (A \cup B))}{P(A \cup B)} = \frac{P(E)}{P(A \cup B)} = \frac{P(A \cup B) - P(A \cap B)}{P(A \cup B)} = \frac{0.4 - 0.1}{0.4} = \frac{3}{4}$.

2.4.5 The answer would remain the same. Distinguishing only three family types does not make them equally likely; (girl, boy) families will occur twice as often as either (boy, boy) or (girl, girl) families.

2.4.6 $P(A \cup B) = 0.8$ and $P(A \cup B) - P(A \cap B) = 0.6$, so $P(A \cap B) = 0.2$.

Also, $P(A|B) = 0.6 = \frac{P(A \cap B)}{P(B)}$, so $P(B) = \frac{0.2}{0.6} = \frac{1}{3}$ and $P(A) = 0.8 + 0.2 - \frac{1}{3} = \frac{2}{3}$.

2.4.7 Let R_i be the event that a red chip is selected on the i th draw, $i = 1, 2$.

Then $P(\text{both are red}) = P(R_1 \cap R_2) = P(R_2 | R_1)P(R_1) = \frac{3}{4} \cdot \frac{1}{2} = \frac{3}{8}$.

2.4.8 $P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{P(A) + P(B) - P(A \cup B)}{P(B)} = \frac{a + b - P(A \cup B)}{b}$.

But $P(A \cup B) \leq 1$, so $P(A|B) \geq \frac{a + b - 1}{b}$.

2.4.9 Let W_i be the event that a white chip is selected on the i th draw, $i = 1, 2$.

Then $P(W_2|W_1) = \frac{P(W_1 \cap W_2)}{P(W_1)}$. If both chips in the urn are white, $P(W_1) = 1$;

if one is white and one is black, $P(W_1) = \frac{1}{2}$.

Since each chip distribution is equally likely, $P(W_1) = 1 \cdot \frac{1}{2} + \frac{1}{2} \cdot \frac{1}{2} = \frac{3}{4}$.

Similarly, $P(W_1 \cap W_2) = 1 \cdot \frac{1}{2} + \frac{1}{4} \cdot \frac{1}{2} = \frac{5}{8}$, so $P(W_2|W_1) = \frac{5/8}{3/4} = \frac{5}{6}$.

2.4.10 $P[(A \cap B) | (A \cup B)^c] = \frac{P[(A \cap B) \cap (A \cup B)^c]}{P[(A \cup B)^c]} = \frac{P(\emptyset)}{P[(A \cup B)^c]} = 0$

2.4.11 (a) $P(A^c \cap B^c) = 1 - P(A \cup B) = 1 - [P(A) + P(B) - P(A \cap B)] = 1 - [0.65 + 0.55 - 0.25] = 0.05$

- (b) $P[(A^C \cap B) \cup (A \cap B^C)] = P(A^C \cap B) + P(A \cap B^C) =$
 $[P(A) - P(A \cap B)] + [P(B) - P(A \cap B)] = [0.65 - 0.25] + [0.55 - 0.25] = 0.70$
- (c) $P(A \cup B) = 0.95$
- (d) $P[(A \cap B)^C] = 1 - P(A \cap B) = 1 - 0.25 = 0.75$
- (e) $P\{[(A^C \cap B) \cup (A \cap B^C)] | A \cup B\} = \frac{P[(A^C \cap B) \cup (A \cap B^C)]}{P(A \cup B)} = 0.70/0.95 = 70/95$
- (f) $P(A \cap B) | A \cup B = P(A \cap B)/P(A \cup B) = 0.25/0.95 = 25/95$
- (g) $P(B|A^C) = P(A^C \cap B)/P(A^C) = [P(B) - P(A \cap B)]/[1 - P(A)] = [0.55 - 0.25]/[1 - 0.65] = 30/35$

- 2.4.12** $P(\text{No. of heads} \geq 2 | \text{No. of heads} \leq 2)$
 $= P(\text{No. of heads} \geq 2 \text{ and No. of heads} \leq 2)/P(\text{No. of heads} \leq 2)$
 $= P(\text{No. of heads} = 2)/P(\text{No. of heads} \leq 2) = (3/8)/(7/8) = 3/7$
- 2.4.13** $P(\text{first die} \geq 4 | \text{sum} = 8) = P(\text{first die} \geq 4 \text{ and sum} = 8)/P(\text{sum} = 8)$
 $= P(\{(4, 4), (5, 3), (6, 2)\})/P(\{(2, 6), (3, 5), (4, 4), (5, 3), (6, 2)\}) = 3/5$
- 2.4.14** There are 4 ways to choose three aces (count which one is left out). There are 48 ways to choose the card that is not an ace, so there are $4 \times 48 = 192$ sets of cards where exactly three are aces. That gives 193 sets where there are at least three aces. The conditional probability is $(1/270,725)/(193/270,725) = 1/193$.
- 2.4.15** First note that $P(A \cup B) = 1 - P[(A \cup B)^C] = 1 - 0.2 = 0.8$.
Then $P(B) = P(A \cup B) - P(A \cap B^C) - P(A \cap B) = 0.8 - 0.3 - 0.1 = 0.5$.
Finally $P(A|B) = P(A \cap B)/P(B) = 0.1/0.5 = 1/5$
- 2.4.16** $P(A|B) = 0.5$ implies $P(A \cap B) = 0.5P(B)$. $P(B|A) = 0.4$ implies $P(A \cap B) = (0.4)P(A)$.
Thus, $0.5P(B) = 0.4P(A)$ or $P(B) = 0.8P(A)$.
Then, $0.9 = P(A) + P(B) = P(A) + 0.8P(A)$ or $P(A) = 0.9/1.8 = 0.5$.
- 2.4.17** $P[(A \cap B)^C] = P[(A \cup B)^C] + P(A \cap B^C) + P(A^C \cap B) = 0.2 + 0.1 + 0.3 = 0.6$
 $P(A \cup B | (A \cap B)^C) = P[(A \cap B^C) \cup (A^C \cap B)]/P((A \cap B)^C) = [0.1 + 0.3]/0.6 = 2/3$
- 2.4.18** $P(\text{sum} \geq 8 | \text{at least one die shows 5})$
 $= P(\text{sum} \geq 8 \text{ and at least one die shows 5})/P(\text{at least one die shows 5})$
 $= P(\{(5, 3), (5, 4), (5, 6), (3, 5), (4, 5), (6, 5), (5, 5)\})/(11/36) = 7/11$
- 2.4.19** $P(\text{Outandout wins} | \text{Australian Doll and Dusty Stake don't win})$
 $= P(\text{Outandout wins and Australian Doll and Dusty Stake don't win})/P(\text{Australian Doll and Dusty Stake don't win}) = 0.20/0.55 = 20/55$
- 2.4.20** Suppose the guard will randomly choose to name Bob or Charley if they are the two to go free. Then the probability the guard will name Bob, for example, is $P(\text{Andy, Bob}) + (1/2)P(\text{Bob, Charley}) = 1/3 + (1/2)(1/3) = 1/2$.
The probability Andy will go free given the guard names Bob is $P(\text{Andy, Bob})/P(\text{Guard names Bob}) = (1/3)/(1/2) = 2/3$. A similar argument holds for the guard naming Charley. Andy's concern is not justified.